

BEFORE SURGERY

1 Day Before Surgery: Start using DROP#1 and DROP #2

1

**Antibiotic
(Tan Top)**



OR



Vigamox

Moxifloxacin/Ofloxacin (generic)

-1 Drop 4 Times a Day

2

**NSAID
(Grey Top)**



OR



Prolensa

Diclofenac (generic)

-If Prolensa, 1 Drop in your operative eye before bed.
-If Diclofenac, then use 2x/day
(Diabetics start 1 week prior to surgery)

- Space eye drops 5 minutes apart.
- Do not touch the tip of the bottles with your finger, to your face, or eye.

MORNING OF SURGERY

- **Do not eat or drink anything 8 hours before your surgery.**
- Expect to be at the surgery facility for 3-4 hours.
- **You MUST have someone take you home after surgery.**
- Unless otherwise instructed by your Doctor, take your usual medications with a small amount of water. **DO NOT** eat or drink anything else.
- Pause any diabetic medication.
- Bring your eye drops + list of medications you are taking to the Surgery Facility.

AFTER SURGERY

1

**Antibiotic
(Tan Top)**



OR



Vigamox

Moxifloxacin/Ofloxacin (generic)

1 Drop 4 Times a Day

2

**NSAID
(Grey Top)**



OR



Prolensa

Diclofenac (generic)

-If Prolensa, 1 Drop in your operative eye before bed.
-If Diclofenac, then use 2x/day

3

**Steroid Drop
(White or
Pink Top)**



OR



Durezol or Inveltys

1 Drop, 2x/day for 2 weeks
Then 1 Drop, 1x/day for 1 week

Prednisolone Acetate 1% (generic)

1 Drop, 4x/day for 2 weeks
Then 1 Drop, 2x/day for 1 week
Then 1 Drop, 1x/day for 1 week

- Wear the shield at bedtime for 1 weeks after surgery on your operative eye. **DO NOT RUB YOUR EYES.**
- No heavy exercise + lifting more than 20 lbs for 2 weeks. Walking is OK.
- No bending with head below the waist for 2 weeks.
- Avoid water in the eyes; No swimming or jacuzzi use for 3 weeks after surgery.
- You may experience blurred vision in your operative eye for a few days or more.
- You may continue to use your artificial drops whenever your eyes dry out.
- **Pause the following glaucoma medications for one month: Latanoprost, Xalatan, Lumigan, and Travatan**
- **Call us ASAP if you have a sudden decrease in vision, flashes, floaters or serious pain with redness.**